

ON LIN'S ARITHMETIC–GEOMETRIC MEANS CONJECTURE FOR SINGULAR VALUES

SUVRIT SRA*

ABSTRACT. Let A and B be positive semidefinite, and let $0 \leq t \leq 1$. Lin (2017) conjectured that

$$\lambda_j((1-t)A + tB) \geq \sigma_j^{1/2}(A^{2(1-t)}B^{2t}), \quad j = 1, \dots, n.$$

The case $t = \frac{1}{2}$ is Drury's theorem, which answered a long-standing question of Bhatia and Kittaneh. Lin remarked that Drury's method does not seem to reach other values of t . We prove his conjecture for every $t \in [0, 1]$. The proof builds on Drury's argument, but replaces his block-matrix step (unique to the midpoint $t = \frac{1}{2}$) by an eigenvalue count based on an inertia identity and a one-variable inequality.

1. INTRODUCTION

Finding matrix analogues of scalar inequalities has long been a theme of matrix analysis. Many works study analogues of the arithmetic–geometric mean inequality; see, e.g., the books (Bhatia, 1997, 2007; Zhan, 2002). Some of the most useful of these inequalities compare singular values $\sigma(\cdot)$ and eigenvalues $\lambda(\cdot)$ of matrices. A fundamental example of such an inequality is

$$(1.1) \quad \sigma_j^{1/2}(AB) \leq \frac{1}{2}\lambda_j(A+B), \quad A, B \succeq 0, \quad j = 1, \dots, n,$$

which was raised as a question by Bhatia and Kittaneh (2000) and proved twelve years later by Drury (2012).

Bhatia and Kittaneh (1990) proved the weaker inequality $\sigma_j(A^{1/2}B^{1/2}) \leq \frac{1}{2}\lambda_j(A+B)$. This inequality was generalized by Ando (1995) to $\sigma_j(A^{1-t}B^t) \leq \lambda_j((1-t)A + tB)$ for $t \in [0, 1]$. Revisiting Drury's proof, Lin (2017, Conjecture 5.2) conjectured the corresponding weighted form of (1.1), and wrote that Drury's method “does not seem to lead to a solution.” To our knowledge no proof has appeared since, and Theorem 1.1 supplies one.

Theorem 1.1 (Lin's conjecture). *Let $A, B \succeq 0$ and let $0 \leq t \leq 1$. Then, for every $j = 1, \dots, n$,*

$$(1.2) \quad \sigma_j^{1/2}(A^{2(1-t)}B^{2t}) \leq \lambda_j((1-t)A + tB).$$

More broadly, both Ando's inequality and Theorem 1.1 are members of the family

$$(1.3) \quad \sigma_j^q(A^{(1-t)/q}B^{t/q}) \leq \lambda_j((1-t)A + tB), \quad j = 1, \dots, n,$$

indexed by $q > 0$. A brief calculation shows that if inequality (1.3) holds for an exponent q , then it also holds for every larger exponent, so the statements become stronger as q decreases. The cases $q \geq 1$ therefore follow from Ando's inequality, while Lin's conjecture is the case $q = 1/2$, outside that range. Theorem 1.1 gives (1.3) for every $q \geq 1/2$, and another brief calculation shows that at $t = 1/2$ the exponent $1/2$ is the best.

* TU Munich, Dept. of Mathematics, School of CIT; Garching, Germany. Email: s.sra@tum.de.

First public release: 8 October 2026, as v1 in the author's preprint repository, <https://github.com/suvrit/preprints/tree/main/preprints/linconj>; every version is kept there with its timestamp.

2. PROOF OF THEOREM 1.1

Proof summary. The proof has four steps. The first two follow Drury's argument (Drury, 2012) expositied in Lin (2017, Sections 2–4). Write $a = 1 - t$ and $b = t$. In Step 1 we use homogeneity of (1.2) to reduce the task to proving $\lambda_j(aA + bB) \geq 1$. Step 2 reduces this task to an eigenvalue inequality for $aA + bB_0$, where $B_0 \preceq B$ is a rank- j matrix. Drury's proof of that inequality works only at $t = 1/2$, so we introduce Steps 3 and 4 to replace it. Step 3 counts eigenvalues using an inertia identity, which reduces the inequality to the sign of a quadratic form on a subspace; Step 4 settles this sign using an elementary scalar inequality.

2.1. Rescaling. Using a and b , inequality (1.2) reads

$$(2.1) \quad \lambda_j(aA + bB) \geq \sigma_j^{1/2}(A^{2a}B^{2b}), \quad j = 1, \dots, n.$$

If $t = 0$ or $t = 1$ the inequality is trivial; so we assume $t \in (0, 1)$. Also, it suffices to consider $A, B \succ 0$ (the $\succeq 0$ cases follow by continuity). We can also assume $a \leq b$ (otherwise flip the roles of A and B and note $\sigma(AB) = \sigma(BA)$), so that $b \geq 1/2$. Fix $j \in \{1, \dots, n\}$. If $\sigma_j(A^{2a}B^{2b}) = 0$, then (2.1) is trivial; otherwise by homogeneity we may scale (2.1) so that $\sigma_j(A^{2a}B^{2b}) = 1$, and reduce the task to proving that $\lambda_j(aA + bB) \geq 1$.

2.2. Rank reduction. By Weyl's monotonicity principle, lowering B in the Löwner order can only lower $\lambda_j(aA + bB)$. Following Drury, we lower B to a matrix B_0 of rank j by keeping only the part of B that produces the j largest singular values of $A^{2a}B^{2b}$, and resetting those singular values to 1.

More precisely, let $T = A^{2a}B^{4b}A^{2a} = (A^{2a}B^{2b})(A^{2a}B^{2b})^*$. Since $\lambda(T) = \sigma^2(A^{2a}B^{2b})$, the rescaling step tells us $\lambda_j(T) = 1$. Choose an orthonormal basis $v_1, \dots, v_n$ with $Tv_k = \lambda_k(T)v_k$, and let E be the orthogonal projection onto $\text{span}(v_1, \dots, v_j)$. Since $T \succeq 0$ and $\lambda_k(T) \geq \lambda_j(T) = 1$ for $k \leq j$,

$$T - E = \sum_{k \leq j} (\lambda_k(T) - 1)v_kv_k^* + \sum_{k > j} \lambda_k(T)v_kv_k^* \succeq 0.$$

Thus, $T \succeq E$. Congruence with A^{-2a} then yields

$$(2.2) \quad B^{4b} = A^{-2a}TA^{-2a} \succeq B_0^{4b} := A^{-2a}EA^{-2a}.$$

Because $b \geq 1/2$, the exponent $1/(4b)$ is at most 1, so by Löwner–Heinz on (2.2) we obtain $B \succeq B_0$. Hence by Weyl $\lambda_j(aA + bB) \geq \lambda_j(aA + bB_0)$.

Why such a choice of B_0 ? The point of this choice is that B_0 is determined by A and by its own support projection P . Let $V = A^{2a}B_0^{2b}$. By (2.2), $VV^* = A^{2a}B_0^{4b}A^{2a} = E$, so V is a partial isometry with final projection E . Since A^{2a} is invertible, $\text{Ker } V = \text{Ker } B_0^{2b} = \text{Ran}(I - P)$, so the initial projection of V is P . In other words,

$$(2.3) \quad B_0^{2b}A^{4a}B_0^{2b} = P.$$

The rank of P is j , because $B_0^{4b} = A^{-2a}EA^{-2a}$ has rank j . Let $S = PA^{4a}P|_{\text{Ran } P}$ be the compression of A^{4a} to $\text{Ran } P$, which is positive definite because A is. Write $B_0 = PXP$ with $X \succ 0$ on $\text{Ran } P$, and let $Y = X^{2b}$. Compressing (2.3) to $\text{Ran } P$ gives $YSY = I_{\text{Ran } P}$, and multiplying on the left and on the right by Y^{-1} gives $S = Y^{-2}$. Since $Y \succ 0$, uniqueness of positive square roots gives $Y = S^{-1/2}$, and therefore $X = S^{-1/(4b)}$. Thus

$$(2.4) \quad B_0 = PS^{-1/(4b)}P.$$

It remains to prove

$$(2.5) \quad \lambda_j(aA + bB_0) \geq 1.$$

Since (2.4) expresses B_0 through A and P alone, we prove (2.5) for every $A \succ 0$ and every orthogonal projection P of rank j , with B_0 defined by (2.4). At $t = 1/2$ the matrix $B_0 = P(PA^2P)^{-1/2}P$ is Drury's matrix B_1 (Lin, 2017, Section 2), and (2.5) is the inequality $\lambda_j(A + B_1) \geq 2$ that Drury

proves by an eigenvalue estimate for 2×2 block matrices. That estimate uses that B_0 is the inverse square root of PA^2P on $\text{Ran } P$, and we see no analogue of it for the fractional power $S^{-1/(4b)}$.

2.3. Eigenvalue count. Inequality (2.5) says that $aA + bB_0 - I$ has at least j nonnegative eigenvalues. This matrix is $H + bB_0$ with $H = aA - I$: a Hermitian matrix, in general indefinite, plus a positive semidefinite perturbation carried by the j -dimensional space $\text{Ran } P$. The following inertia identity transfers such an eigenvalue count to a matrix that acts only on the space carrying the perturbation. It is the finite-dimensional form of the Birman–Schwinger principle (Schwinger, 1961).

Lemma 2.1 (Inertia identity). *Let $H = H^*$ be invertible on a finite-dimensional Hilbert space $\mathcal{H}$. Let $J: \mathcal{P} \rightarrow \mathcal{H}$ be an isometric inclusion, and let $W \succ 0$ on $\mathcal{P}$. Set*

$$M = H + JWJ^*, \quad G = W^{-1} + J^*H^{-1}J.$$

Then

$$n_{\geq 0}(M) = n_+(H) + n_{\leq 0}(G),$$

where $n_{\geq 0}$ counts eigenvalues ≥ 0 , n_+ counts eigenvalues > 0 , and $n_{\leq 0}$ counts eigenvalues ≤ 0 , each with multiplicity.

Proof. Consider the Hermitian matrix

$$\mathcal{L} = \begin{pmatrix} H & J \\ J^* & -W^{-1} \end{pmatrix} \quad \text{on } \mathcal{H} \oplus \mathcal{P}.$$

The Schur complements in $\mathcal{L}$ of its blocks $-W^{-1}$ and H are

$$H + JWJ^* = M \quad \text{and} \quad -W^{-1} - J^*H^{-1}J = -G.$$

By inertia additivity for Schur complements (Haynsworth, 1968), $\mathcal{L}$ therefore has the inertia of both $M \oplus (-W^{-1})$ and $H \oplus (-G)$. As $-W^{-1} \prec 0$ and H is invertible, counting nonnegative eigenvalues gives $n_{\geq 0}(M) = n_{\geq 0}(\mathcal{L}) = n_+(H) + n_{\leq 0}(G)$. $\square$

To apply Lemma 2.1, $H = aA - I$ must be invertible, that is, $1/a$ must not be an eigenvalue of A . By continuity we may assume so: for fixed P , the left side of (2.5) depends continuously on $A \succ 0$, and $1/a$ is an eigenvalue of $A + \varepsilon I$ for only finitely many $\varepsilon > 0$.

Let m be the number of eigenvalues of A greater than $1/a$, so that $n_+(H) = m$. Apply Lemma 2.1 with $\mathcal{H} = \mathbb{C}^n$, $\mathcal{P} = \text{Ran } P$, J the inclusion map, and $W = bS^{-1/(4b)} \succ 0$. Then $JWJ^* = bB_0$, so $M = aA + bB_0 - I$, and since $(aA - I)^{-1} = -(I - aA)^{-1}$,

$$(2.6) \quad n_{\geq 0}(aA + bB_0 - I) = m + n_{\leq 0}(G), \quad G = b^{-1}S^{1/(4b)} - P(I - aA)^{-1}P|_{\text{Ran } P}.$$

Thus (2.5) holds as soon as G has at least $j - m$ nonpositive eigenvalues. It is enough to find a subspace $U \subseteq \text{Ran } P$ with $\dim U \geq j - m$ on which the quadratic form of G is nonpositive. Indeed, for such a subspace $n_{\leq 0}(G) \geq \dim U$: otherwise the span N of the eigenvectors of G for eigenvalues ≤ 0 would have $\dim N < \dim U$, so U would contain a nonzero vector $v \perp N$, and $\langle Gv, v \rangle > 0$.

The two terms of G indicate where to look. The first is positive definite. The second, $-P(I - aA)^{-1}P$, is negative on vectors built from eigenvectors of A with eigenvalues $x < 1/a$, on which $(I - aA)^{-1}$ acts by the positive numbers $1/(1 - ax)$. Let $\mathcal{F}$ be the span of the eigenvectors of A with eigenvalues in $[0, 1/a)$, and let $U = \text{Ran } P \cap \mathcal{F}$. Because $1/a$ is not an eigenvalue of A , $\mathcal{F}$ has dimension $n - m$, and since $\text{Ran } P$ has dimension j ,

$$\dim U \geq j + (n - m) - n = j - m.$$

It therefore remains to prove

$$(2.7) \quad \langle Gu, u \rangle \leq 0 \quad \text{for every } u \in U.$$

2.4. Scalar inequality. By homogeneity it suffices to prove (2.7) for a unit vector $u \in U$. Let μ be the spectral measure of A associated with u , that is, the probability measure on $\mathbb{R}$ with $\langle f(A)u, u \rangle = \int f d\mu$ for every function f on the spectrum of A . Since $u \in \mathcal{F}$, the measure μ is supported in $[0, 1/a)$. Since $u \in \text{Ran } P$,

$$\langle Su, u \rangle = \langle A^{4a}u, u \rangle = \int x^{4a} d\mu(x) \quad \text{and} \quad \langle P(I - aA)^{-1}Pu, u \rangle = \int \frac{d\mu(x)}{1-ax}.$$

Put $\gamma = 1/(4b)$; because $b \geq 1/2$, we have $0 < \gamma \leq 1$. Concavity of $x \mapsto x^\gamma$ and the spectral measure of S associated with u give $\langle S^\gamma u, u \rangle \leq \langle Su, u \rangle^\gamma$. By (2.6), therefore,

$$(2.8) \quad \langle Gu, u \rangle \leq b^{-1} \left(\int x^{4a} d\mu(x) \right)^\gamma - \int \frac{d\mu(x)}{1-ax}.$$

So (2.7) follows once the right side of (2.8) is shown to be nonpositive for every probability measure μ on $[0, 1/a)$. Lemma 2.2 proves this bound by integrating an inequality in one real variable; its proof is in Appendix A.

Lemma 2.2 (Scalar inequality). *Let $0 < a \leq b$, $a + b = 1$. For every $0 \leq x < 1/a$,*

$$(2.9) \quad x^{4a} \leq 1 + 4b^2 \left(\frac{1}{1-ax} - \frac{1}{b} \right).$$

Consequently, if μ is a probability measure on $[0, 1/a)$, then

$$(2.10) \quad \left(\int x^{4a} d\mu(x) \right)^{1/(4b)} \leq b \int \frac{1}{1-ax} d\mu(x).$$

By (2.10), the right side of (2.8) is nonpositive. This proves (2.7), and with it Theorem 1.1. $\square$

AI USAGE

The results in this paper were developed with an AI assistant (OpenAI's GPT Pro). None of the claims were obtained zero-shot: they emerged through sustained back-and-forth discussion, including failed attempts, counterexamples, audits, and repairs.

ACKNOWLEDGEMENTS

The author (SS) thanks the Alexander von Humboldt (AvH) Foundation for their generous support via a Humboldt Professorship in AI.

REFERENCES

- Tsuyoshi Ando. Matrix Young inequalities. In *Operator Theory in Function Spaces and Banach Lattices*, volume 75 of *Operator Theory: Advances and Applications*, pages 33–38. Birkhäuser, Basel, 1995. doi: 10.1007/978-3-0348-9076-2_5.
- Rajendra Bhatia. *Matrix Analysis*. Graduate Texts in Mathematics 169. Springer, New York, 1997. doi: 10.1007/978-1-4612-0653-8.
- Rajendra Bhatia. *Positive Definite Matrices*. Princeton Series in Applied Mathematics. Princeton University Press, Princeton, NJ, 2007. doi: 10.1515/9781400827787.
- Rajendra Bhatia and Fuad Kittaneh. On the singular values of a product of operators. *SIAM Journal on Matrix Analysis and Applications*, 11(2):272–277, 1990. doi: 10.1137/0611018.
- Rajendra Bhatia and Fuad Kittaneh. Notes on matrix arithmetic–geometric mean inequalities. *Linear Algebra and its Applications*, 308(1-3):203–211, 2000. doi: 10.1016/S0024-3795(00)00048-3.
- Stephen W. Drury. On a question of Bhatia and Kittaneh. *Linear Algebra and its Applications*, 437(7):1955–1960, 2012. doi: 10.1016/j.laa.2012.04.040.
- Emilie V. Haynsworth. Determination of the inertia of a partitioned Hermitian matrix. *Linear Algebra and its Applications*, 1(1):73–81, 1968. doi: 10.1016/0024-3795(68)90050-5.

- Minghua Lin. On Drury's solution of Bhatia and Kittaneh's question. *Linear Algebra and its Applications*, 528:33–39, 2017. doi: 10.1016/j.laa.2016.02.025.
- Julian Schwinger. On the bound states of a given potential. *Proceedings of the National Academy of Sciences*, 47(1):122–129, 1961. doi: 10.1073/pnas.47.1.122.
- Xingzhi Zhan. *Matrix Inequalities*. Lecture Notes in Mathematics 1790. Springer, Berlin, 2002. doi: 10.1007/b83956.

APPENDIX A. PROOF OF THE SCALAR INEQUALITY

Equality holds in (2.9) at $x = 1$, where both sides equal 1 and both have derivative $4a$, so the two sides are tangent there.

Proof. Put $y = (1 - ax)^{-1}$, so that $y \geq 1$ and $ax = 1 - y^{-1}$. Then (2.9) says $g(y) \geq 0$ for $y \geq 1$, where $g(y) = 1 + 4b^2(y - b^{-1}) - a^{-4a}(1 - y^{-1})^{4a}$. Since $1 - b = a$, we have $g(b^{-1}) = 0$, and $g'(y) = 4b^2 - 4a^{1-4a}(1 - y^{-1})^{4a-1}y^{-2}$ also vanishes at b^{-1} ; moreover $g(1) = (1 - 2a)^2 \geq 0$. For $y > 1$,

$$g''(y) = 4a^{1-4a} \frac{(1 - y^{-1})^{4a}(2y - 1 - 4a)}{y^2(y - 1)^2},$$

so g is concave on $[1, y_*]$ and convex on $[y_*, \infty)$, where $y_* = (1 + 4a)/2$ (convex throughout if $y_* \leq 1$). Since $2 - (1 + 4a)(1 - a) = 4a^2 - 3a + 1 > 0$, we have $y_* < b^{-1}$. On the convex part g has a double zero at b^{-1} , so $g \geq 0$ there, in particular at y_* ; on the concave part g lies above the chord joining $g(1) \geq 0$ and $g(y_*) \geq 0$. This proves (2.9).

Integrating (2.9) against μ gives $\int x^{4a} d\mu \leq 1 + 4b^2(\bar{y} - b^{-1})$ with $\bar{y} = \int (1 - ax)^{-1} d\mu$. The right side is the tangent line at b^{-1} of $y \mapsto (by)^{4b}$, which is convex since $4b \geq 1$. Hence $\int x^{4a} d\mu \leq (b\bar{y})^{4b}$, and taking the power $1/(4b)$ gives (2.10). $\square$